



Contents lists available at ScienceDirect

Journal of Colloid and Interface Science

www.elsevier.com/locate/jcis



Short Communication

Comments on “Adsorption of direct dyes from aqueous solutions by carbon nanotubes: Determination of equilibrium, kinetics and thermodynamics parameters”

Yuh-Shan Ho*

Department of Environmental Sciences, College of Environmental Science and Engineering, Peking University, Beijing, 100871, People's Republic of China

ARTICLE INFO

Article history:

Received 10 December 2008

Accepted 11 January 2009

Available online 12 February 2009

Keywords:

Initial adsorption rate

Pseudo-second-order kinetic model

Intraparticle diffusion model

Quotation error

ABSTRACT

Two most suggested papers for pseudo-second-order kinetic model were published in 1984 and 1995 by Blanchard et al. and Ho, respectively. Blanchard et al. noted the overall exchange reaction of NH_4^+ ions fixed in zeolite by divalent metallic ions in the solution using a second-order kinetic model. Ho used the pseudo-second-order kinetic model to the copper ion/peat adsorption system. The adsorption involved chemical bonding and cation exchange. In this comment citation error and quotation error were pointed.

© 2009 Elsevier Inc. All rights reserved.

Recently, Kuo et al. [1] published the paper noted above. In Section 3.2.1, Effects of dye concentration, the authors noted a pseudo-second-order model, initial adsorption rate h , and intraparticle diffusion model. In the case of the pseudo-second-order model the authors cited two secondary materials as references [2,3]. In fact, the pseudo-second-order kinetic expression for the adsorption systems of divalent metal ions using sphagnum moss peat has been presented by Ho [4] and this kinetic expression has also been published in 1996 [5]. At the same time Ho has presented a definition for the initial adsorption rate from the pseudo-second-order model. A modified model has been made in the following years because a mistake was included in the paper published in 1996 [6–8]. However, in the case of initial adsorption rate, the authors noted that “the initial adsorption rate h (mg/g min) can be determined by using the equation $h = k_2 q_e^2$ ” without any citations. In the case of the intraparticle diffusion model, the authors cited the same two secondary materials as references [2,3]. A quotation error of the intraparticle diffusion model was contained in

both references. The equation presented in the paper could not be found in the original paper of Weber and Morris [9].

Accuracy of quotations and citations is very important for the transmission of scientific knowledge. I suggest that Kuo et al. cite the original or the most frequently cited papers for the kinetic models to have more accuracy and details of information about kinetic expression.

References

- [1] C.Y. Kuo, C.H. Wu, J.Y. Wu, J. Colloid Interface Sci. 327 (2008) 308.
- [2] C.H. Wu, J. Hazard. Mater. 144 (2007) 93.
- [3] I.D. Mall, V.C. Srivastava, N.K. Agarwal, Dyes Pigments 69 (2006) 210.
- [4] Y.S. Ho, Ph.D. thesis, University of Birmingham, Birmingham, UK, 1995.
- [5] Y.S. Ho, D.A.J. Wase, C.F. Forster, Environ. Technol. 17 (1996) 71.
- [6] Y.S. Ho, G. McKay, Chem. Eng. J. 70 (1998) 115.
- [7] Y.S. Ho, G. McKay, Water Res. 34 (2000) 735.
- [8] Y.S. Ho, J. Hazard. Mater. 136 (2006) 681.
- [9] W.J. Weber Jr., J.C. Morris, J. Sanit. Eng. Div. ASCE 89 (1963) 31.

DOI of original article: 10.1016/j.jcis.2008.08.038.

* Fax: +86 10 62751923.

E-mail address: dr_ysho@hotmail.com.