


Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.




This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

Available at [www.sciencedirect.com](http://www.sciencedirect.com)journal homepage: [www.elsevier.com/locate/watres](http://www.elsevier.com/locate/watres)

## Comment

# Comment on “Adsorption mechanism of selenate and selenite on the binary oxide systems” by Y.T. Chan et al. [Water Research 43 (2009) 4412-4420]

Yuh-Shan Ho\*

Department of Environmental Sciences, College of Environmental Science and Engineering, Peking University, Beijing 100871, People's Republic of China

## ARTICLE INFO

### Article history:

Received 4 November 2009

Accepted 14 April 2010

Available online 24 April 2010

Recently, Chan et al. (2009) published the paper entitled above. In section of 2.4. Adsorption kinetics, the authors noticed “the initial sorption rate” with Eq. (4) without any citations. In fact, Ho has presented a definition for the initial adsorption rate from the parameters of pseudo-second-order model (Ho, 1995; Ho et al., 1996). A modified initial adsorption rate equation has also been made in the following years because a mistake was included in the paper published in 1996 (Ho and McKay, 1998; Ho, 2006). In addition, authors cited Saha et al. (2004) for pseudo-second-order kinetic model. In fact there is nothing related the model in the reference.

Accuracy of citations and quotations are very important for the transmission of scientific knowledge. I suggest that Chan et al. cite the original or the most frequently cited papers for the initial adsorption rate to have more accuracy and details of information about its expression.

## REFERENCES

- Chan, Y.T., Kuan, W.H., Chen, T.Y., Wang, M.K., 2009. Adsorption mechanism of selenate and selenite on the binary oxide systems. *Water Res.* 43, 4412–4420.
- Ho, Y.S., 1995. Adsorption of heavy metals from waste streams by peat. Ph.D. thesis, The University of Birmingham, Birmingham, U.K.
- Ho, Y.S., 2006. Review of second-order models for adsorption systems. *J. Hazard. Mater.* 136, 681–689.
- Ho, Y.S., McKay, G., 1998. Sorption of dye from aqueous solution by peat. *Chem. Eng. J.* 70, 115–124.
- Ho, Y.S., Wase, D.A.J., Forster, C.F., 1996. Kinetic studies of competitive heavy metal adsorption by sphagnum moss peat. *Environ. Technol.* 17, 71–77.
- Saha, U.K., Liu, C., Kozak, L.M., Huang, P.M., 2004. Kinetics of selenite adsorption on hydroxyaluminum- and hydroxyaluminosilicate–montmorillonite complexes. *Soil Sci. Soc. Am. J.* 68 (4), 1197–1209.

DOI of original article: 10.1016/j.watres.2009.06.056.

\* Tel./fax: +86 10 62751923.

E-mail address: [dr\\_ysho@hotmail.com](mailto:dr_ysho@hotmail.com)

0043-1354/\$ – see front matter © 2010 Elsevier Ltd. All rights reserved.

doi:10.1016/j.watres.2010.04.013